

By Seth A. Berkowitz, Jean Terranova, Caterina Hill, Toyin Ajayi, Todd Linsky, Lori W. Tishler, and Darren A. DeWalt

Meal Delivery Programs Reduce The Use Of Costly Health Care In Dually Eligible Medicare And Medicaid Beneficiaries

DOI: 10.1377/hlthaff.2017.0999
 HEALTH AFFAIRS 37,
 NO. 4 (2018): 535–542
 ©2018 Project HOPE—
 The People-to-People Health
 Foundation, Inc.

ABSTRACT Delivering food to nutritionally vulnerable patients is important for addressing these patients' social determinants of health. However, it is not known whether food delivery programs can reduce the use of costly health services and decrease medical spending among these patients. We sought to determine whether home delivery of either medically tailored meals or nontailored food reduces the use of selected health care services and medical spending in a sample of adults dually eligible for Medicare and Medicaid. Compared with matched nonparticipants, participants had fewer emergency department visits in both the medically tailored meal program and the nontailored food program. Participants in the medically tailored meal program also had fewer inpatient admissions and lower medical spending. Participation in the nontailored food program was not associated with fewer inpatient admissions but was associated with lower medical spending. These findings suggest the potential for meal delivery programs to reduce the use of costly health care and decrease spending for vulnerable patients.

Seth A. Berkowitz (seth_berkowitz@med.unc.edu) is an assistant professor of medicine in the Division of General Medicine and Clinical Epidemiology, School of Medicine, University of North Carolina at Chapel Hill. During the time of the study, he was an assistant professor of medicine at Massachusetts General Hospital/Harvard Medical School, in Boston.

Jean Terranova is director of food and health policy at Community Servings, in Jamaica Plain, Massachusetts.

Caterina Hill is director of research and evaluation at Commonwealth Care Alliance, in Boston.

Toyin Ajayi is chief health officer of Sidewalk Labs Care Lab, in New York City.

Todd Linsky is senior business intelligence informatics analyst at Commonwealth Care Alliance.

Lori W. Tishler is vice president of medical affairs at Commonwealth Care Alliance.

Darren A. DeWalt is the John Randolph and Helen Barnes Chambliss Distinguished Professor of Medicine, Division of General Medicine and Clinical Epidemiology, School of Medicine, University of North Carolina at Chapel Hill.

Social determinants of health, such as lack of access to nutritious food, are recognized as factors associated with high costs of health care.^{1–3} Approximately 13 percent of US households report food insecurity, meaning that they lack “consistent, dependable access to enough food for active, healthy living.”⁴ Food insecurity is associated with poor health and increased use of “big-ticket” health services, such as emergency department (ED) visits and inpatient admissions.^{5–12} Proposed mechanisms of this association include poor dietary quality leading to increased disease complications, making trade-offs between food and medication purchases that impair chronic disease management, and increased stress that worsens mental health.¹³ Perhaps for these reasons, food insecurity is associated with \$77 billion in excess health care expenditures each year.¹⁴

Lack of access to nutritious food may be a particular problem for people who are dually eligible for Medicare and Medicaid.¹⁵ These individuals qualify for Medicaid on the basis of low income and for Medicare through either adjudicated disability or age. Vulnerability from both medical issues and poverty that dually eligible patients face has led to support for services, such as meal programs, that were uncommon in fee-for-service systems.^{3,16} However, it is not yet known whether meal programs improve vulnerable patients’ health care use.

In this study we sought to determine the impact of two types of home-delivered meal interventions, which had different program costs and target populations, on health care use and spending. Because there is an association between food insecurity and greater use of the ED, we hypothesized that the interventions would reduce participants’ ED visits.^{5–8} Second-

arily, we hypothesized that participation in one of the meal programs would reduce the use of other big-ticket services (inpatient admissions and emergency transportation, such as transportation by ambulance), along with associated medical spending.

Study Data And Methods

STUDY POPULATION AND SAMPLE SELECTION The study sample was drawn from members of Commonwealth Care Alliance (CCA), and the primary data source for this study was CCA health care claims. CCA is a not-for-profit community-based health plan that manages and administers care for adults (people older than age twenty-one) who are dually eligible for Medicaid and Medicare.¹⁷ CCA uses capitated payments to provide enhanced primary care and care coordination programs. All CCA members with at least six months of continuous enrollment in one of the two meal delivery programs in the period January 1, 2014–January 1, 2016, were eligible for this study. We randomly selected CCA members who, during the same time period, did not receive either meal program, as a control group. For participants, the index date was the date they began receiving the meals; for the control group, it was a randomly selected date. To have sufficient data for matching, we required twelve months of enrollment in CCA before the index date (see online appendix exhibit A1).¹⁸ Participants were followed until either the end of their membership in CCA or the end of the study period (June 30, 2016). Participants who stopped their meal program were analyzed as part of their meal program group.

The Partners Human Research Committee provided human subjects approval for this project.

MEAL DELIVERY PROGRAMS We studied two meal programs. The first was a medically tailored meals program that provided meals customized to participants' medical needs. Each week the program delivered to participants' homes five days' worth of lunches, dinners, and snacks. A registered dietitian tailored the meals to participants' medical needs across seventeen dietary tracks (for example, diabetes, renal, soft, and so on), with combinations of up to three tracks permitted. The second was a Meals on Wheels-type nontailored food program. It also delivered nutritious meals but did not tailor them to participants' medical needs. It provided five days' worth of prepared lunches and dinners each week, usually delivered daily.

Enrollment in either program was determined by an authorizing clinician. Authorization required a determination that the member was

at nutritional risk (for example, he or she had significant weight change, food scarcity, or medical issues that required a specific diet). There was no specific guidance for clinicians as to whether the member should receive the medically tailored meal or the nontailored food program.

OUTCOMES The primary outcome was ED visits (regardless of whether the patient was admitted as an inpatient or discharged home), obtained through health care claims (January 1, 2014, to June 30, 2016). Secondary outcomes included inpatient admissions and use of emergency transportation—a high-cost component of care in this population. In addition, we examined medical spending (expressed as inflation-adjusted 2016 dollars), using claims for five service categories where nutrition programs might plausibly affect expenditure:¹³ inpatient, outpatient, ED, pharmacy, and emergency transportation.

DATA EXTRACTION From CCA data, we extracted information on age, race/ethnicity, sex, primary language (English or non-English), type of CCA plan (One Care or Senior Care Options), and the start and end (if any) of CCA enrollment. We examined a risk score (from the Centers for Medicare and Medicaid Services' Hierarchical Condition Category risk-adjustment model),¹⁹ and medication classes for prescriptions filled in the prior year. We used *International Classification of Diseases*, Ninth Revision (ICD-9), and *International Statistical Classification of Diseases and Related Health Problems*, Tenth Revision (ICD-10), diagnosis codes in claims data to create a comorbidity index (range: -2 to 26) following the method of Joshua Gagne and colleagues.^{20,21} In addition, we used information about participants' area of residence, as indicated by their ZIP code, based on data from the Census Bureau's American Community Survey²² and from the Dartmouth Atlas of Health Care.²³ (For more details, see the "Data Extraction" section of the appendix).¹⁸

STATISTICAL ANALYSIS Participation in the two food programs did not occur at random, which is a major source of confounding in this study. To account for this, we used a nonparametric approach called coarsened exact matching^{24–27} (see the "Coarsened Exact Matching" section of the appendix).¹⁸ This approach seeks to balance relevant sociodemographic, clinical, and pre-intervention health care characteristics that would lead to treatment between the treated and untreated groups, using only pre-intervention data. The approach leads to matched groups that are similar not only in their mean values of covariates, but also across the entire distribution of values (which is important for analyzing health care use, as a small number of cases can contribute a large portion of the total outcome). We

This study extends our knowledge of the effects of providing home-delivered meals on health outcomes.

created two cohorts using coarsened exact matching—one of people who received the medically tailored meals program and similar participants who received no meal intervention, and the other of people who received the nontailored food program and similar participants who received no meal intervention. Owing to the relatively small sample size and substantial differences between those who received the two types of programs, we were unable to compare the two intervention groups with each other.

After creating the matched cohorts, to account for any imbalance between the groups still present after matching, we conducted regression-adjusted analyses using generalized linear models. For “count” outcomes, we used a generalized linear model with a negative binomial distribution and log link. The models included demographic and clinical variables along with medical spending in the twelve months before the index date (to account for regression to the mean—an analysis-of-covariance approach)²⁸ and the index date (to account for secular trends). To analyze monthly post-intervention medical spending, we used a generalized linear model with a gamma distribution and a log link.²⁹ We express differences in post-intervention medical spending between the groups in terms of both that spending itself (the gross difference) and spending net of intervention costs (the net difference). Intervention costs were calculated by summing the paid claims for the services and dividing by the participant’s follow-up time (number of months) to calculate an average per person monthly cost.

We conducted several sensitivity analyses to check the robustness of the results. First, we sought to determine whether differences could be due to a triggering event, such as an inpatient admission, and we conducted analyses in a sample that matched based on these events (see the “Statistical Analysis” section of the appendix).¹⁸ Next, we ran the same models on the entire (unmatched) sample. After that, we forced a very close match on pre-intervention medical spending, with less emphasis on other factors, and

then conducted gamma regression analyses as described above. Finally, to check whether changes might have been related to other social support services (for example, personal care assistants or home health aides), we examined the use of these services in the period sixty days before and sixty days after the index date and during the entire post-index period.

All models used robust confidence intervals for inference. Analyses were conducted in SAS, version 9.4, and Stata/SE, version 14.2.

LIMITATIONS The findings of this study should be interpreted considering several limitations. First, as noted above, entry into the meal delivery programs was not randomized. Though we accounted for measured factors that could influence program entry, we could not exclude unmeasured confounding—for example, by stigma associated with program participation.

Second, the study sample was drawn from a population of people dually eligible for Medicare and Medicaid who were participating in a health insurance program specifically designed for people with high levels of medical and social complexity. These participants were more diverse than state averages and highly concentrated in urban areas. It is unclear how these results might generalize to different populations.

Third, we did not have data on who was offered, but declined, participation in either program.

Fourth, we present our results using an intention-to-treat analysis strategy. However, this might have biased our results to the null, since including those who discontinued the intervention could have diluted the association between the intervention and changes in utilization.

Finally, we could not assess how concurrent interventions that did not result in health care claims might have affected results.

Study Results

For the analyses of the medically tailored meals program, we included 133 participants who received the meals and 1,002 matched controls. For the analyses of the nontailored food program, we included 624 participants who received the intervention and 1,318 matched controls. In these cohorts, demographic characteristics of the intervention and control groups were generally similar, both for mean values and across the distribution of covariates (see exhibit 1 and appendix exhibits A2a and A2b).¹⁸

Very few patients were lost to follow-up: 99.5 percent (1,129 of 1,135) of the participants in the medically tailored meals program cohort and 96.6 percent (1,875 of 1,942) of those in the nontailored food program cohort were enrolled

EXHIBIT 1

Sociodemographic and clinical characteristics of participants in the Commonwealth Care Alliance medically tailored meals and nontailored food programs and matched controls

	Medically tailored meals program		Nontailored food program	
	Control (n = 1,002)	Intervention (n = 133)	Control (n = 1,318)	Intervention (n = 624)
Mean age, years (SD)	57.9 (5.4)	57.4 (8.4)	73.1 (5.9)	73.5 (7.5)
Female (%)	53.49	55.64	63.78	60.74
Race/ethnicity (%)				
Non-Hispanic white	35.22	37.59	12.50	12.50
Non-Hispanic black	13.78	20.30	0.80	0.80
Hispanic	8.27	8.27	28.53	28.53
Asian, other, multiple, or declined to answer	42.73	33.84	58.17	58.17
Non-English primary language (%)	34.98	27.07	46.86	52.88
Insurance product (%)				
One Care	78.04	81.95	22.55	20.03
Senior Care Options	21.96	18.05	77.45	79.97
Spending in 12 months before intervention				
Mean	\$5,475	\$11,251	\$5,095	\$5,446
SD	3,849	8,553	3,887	5,619
Risk score ^a				
Mean	1.42	1.40	1.54	1.53
SD	0.44	0.69	0.43	0.54
Comorbidity index ^b				
Mean	0.26	0.26	0.17	0.17
SD	0.25	0.39	0.26	0.32
Rural ZIP Code Tabulation Area (%)				
Mean	0.24	0.00	0.38	0.48
SD	0.45	0.00	0.98	1.32
Households in ZCTA living in poverty (%)				
Mean	16.06	16.26	17.96	17.86
SD	3.24	5.14	5.38	6.66
Medicare beneficiaries in ZIP code with primary care provider visit in past 12 months (%)				
Mean	80.50	79.48	80.97	80.97
SD	1.00	1.06	1.22	1.52
Prescribed medications (%)				
Insulin	10.24	18.80	13.71	15.71
Antihypertensive	52.62	59.40	68.11	70.35
Other cardiovascular disease medication	26.14	29.32	40.22	40.22
Anticoagulant	4.42	4.51	6.65	7.05
Proton pump Inhibitor	30.39	48.12	40.01	43.27
Antiretroviral	0.75	0.75	0.89	1.12
Phosphate binder	0.11	0.00	0.75	0.32
Inhaler	15.18	30.08	17.69	19.55
Oral steroids	15.09	25.56	16.37	14.90
Antibiotics	33.22	45.11	36.75	30.61

SOURCE Authors' analysis of data from Commonwealth Care Alliance. **NOTES** Multivariate imbalance was assessed using the L1 statistic (0.97 in the medically tailored meal group and 0.96 in the nontailored food group after matching). Detailed information on balance is in appendix exhibits A2a and A2b (see note 18 in text). SD is standard deviation. ^aRisk score as derived from the CMS-HCC Risk Adjustment Model (see note 19 in text). The range in the sample was 0.20–12.03, with higher values indicating worse health. ^bScore as derived from the Gagne comorbidity index (see note 20 in text). The range in the sample is -2–23, with higher scores indicating worse health.

in CCA through the end of the study period (details of meal program participation are in appendix exhibit A3).¹⁸ Owing to a later start within the study period for the medically tailored meals recipients, average post-index follow-up

time there was 19.1 (standard deviation: 5.8) months for those who received meals and 23.0 (SD: 2.0) months for their controls. Average post-index follow-up time in the nontailored food program cohort was 23.6 (SD: 5.1) months

for those who received meals and 24.3 (SD: 2.6) months for their controls. In the analyses of the medically tailored meals program, mean per person counts of ED, inpatient, and emergency transportation use were 1.8, 0.3, and 1.2, respectively. In the analyses of the nontailored food program, mean per person counts were 1.3, 0.4, and 1.0, respectively.

In negative binomial regression analyses, participation in the medically tailored meals program was associated with fewer ED visits (adjusted incidence rate ratio: 0.30; 95% confidence interval: 0.20, 0.45), inpatient admissions (aIRR: 0.48; 95% CI: 0.26, 0.90), and uses of emergency transportation (aIRR: 0.28; 95% CI: 0.16, 0.51) (exhibit 2, appendix exhibits A4–A7).¹⁸ Participation in the nontailored food program was associated with fewer ED visits (aIRR: 0.56; 95% CI: 0.47, 0.68) and uses of emergency transportation (aIRR: 0.62; 95% CI: 0.49, 0.78), but not with fewer inpatient admissions (aIRR: 0.88; 95% CI: 0.69, 1.11) (the full models are in appendix exhibits A8–A10).¹⁸

In gamma regression models that examined average monthly medical expenditures after the index date, participation in the medically tailored meals program was associated with lower medical spending, compared to not participating (difference: -\$570; 95% CI: -931, -208) (exhibit 3, appendix exhibit A11).¹⁸ Participation in the nontailored food program was also associated with lower medical spending (difference:

-\$156; 95% CI: -308, -5) (appendix exhibit A12).¹⁸ The average monthly program costs per participant were \$350 for the medically tailored meals program and \$146 for the nontailored food program (data not shown). Subtracting the program costs from the estimated savings yielded a net savings of \$220 for the medically tailored meals program and \$10 for the nontailored food program.

We found that the results of our main analyses were robust across a number of sensitivity analyses. As an illustration, matching based on a triggering event rather than pre-intervention medical spending did not substantially alter the results (exhibit 4). Other sensitivity analyses included using the entire (unmatched) cohort, in which results for the medically tailored meals program were qualitatively unchanged and results for the nontailored food program results were also similar—except that the reduction in medical spending was no longer significant ($p = 0.05$), and reductions in inpatient admissions were significant ($p = 0.001$). Sensitivity analyses that forced a very close match on costs again favored receipt of medically tailored meals (difference in cost: -\$706; $p < 0.0001$) (appendix exhibits A13 and A14).¹⁸ We did not find significant differences in the use of, or costs associated with, other supportive programs (such as home health aide or personal care assistant services) between either intervention group and its matched controls (appendix exhibit A15).¹⁸ Ad-

EXHIBIT 2

Estimated absolute and relative changes in use of selected health care services, by intervention

	Mean event count					
	Intervention group		Matched control group		Incidence rate ratio	(95% CI)
	Number	(95% CI)	Number	(95% CI)		
EMERGENCY DEPARTMENT VISITS						
Medically tailored meals program	0.63	(0.39, 0.88)	2.10****	(1.73, 2.47)	0.30****	(0.20, 0.45)
Nontailored food program	0.90	(0.74, 1.06)	1.59****	(1.31, 1.88)	0.56****	(0.47, 0.68)
INPATIENT ADMISSIONS						
Medically tailored meals program	0.27	(0.08, 0.46)	0.56**	(0.34, 0.78)	0.48**	(0.26, 0.90)
Nontailored food program	0.43	(0.33, 0.53)	0.49	(0.38, 0.60)	0.88	(0.69, 1.11)
EMERGENCY TRANSPORTATION EVENTS						
Medically tailored meals program	0.46	(0.19, 0.72)	1.60****	(1.12, 2.09)	0.28	(0.16, 0.51)
Nontailored food program	1.06	(0.73, 1.39)	1.70****	(1.15, 2.24)	0.62****	(0.49, 0.78)

SOURCE Authors' analysis of data from Commonwealth Care Alliance (CCA). **NOTES** The exhibit presents estimated absolute number of events ("count"), absolute difference in number of events between groups ("difference"), and relative differences (incidence rate ratio), with associated confidence intervals (CIs) and p values. Estimates from negative binomial models adjusted for receipt of medically tailored meals program, index year, spending in 12 months before intervention, risk score, comorbidity index, CCA enrollment year, insurance product, age, age squared, percent of ZIP Code Tabulation Area (ZCTA) that was rural, percent of households in ZCTA living in poverty, non-Hispanic white, non-Hispanic black, Hispanic, female, prescribed medication classes before the intervention (see exhibit 1), percent of Medicare beneficiaries in ZIP code with primary care provider visit in past twelve months, English as primary language, and follow-up time. The count represents the estimated difference over the follow-up period. p values for comparison of intervention with matched controls. ** $p < 0.05$ **** $p < 0.001$

EXHIBIT 3

Estimated average monthly medical spending per person, by intervention and in control groups

	Intervention group	Matched control group	Gross difference	Net difference
Medically tailored meals program	\$ 843	\$1,413	-\$570***	-\$220
Nontailored food program	\$1,007	\$1,163	-\$156**	-\$10

SOURCE Authors' analysis of data from Commonwealth Care Alliance. **NOTES** Spending is in inflation-adjusted 2016 dollars. Estimates from gamma regression models adjusted for the factors listed in the notes to exhibit 2. Gross difference represents the estimated difference in health care spending by intervention status. Net difference represents the estimated difference in health care expenditures, accounting for the cost of the intervention. *p* values test the null hypothesis that the difference in gross spending between intervention and matched controls is equal to zero. ***p* < 0.05 ****p* < 0.01

justing for supportive services use and cost did not substantially alter the estimates of cost savings, which remained in favor of the intervention programs (appendix exhibit A16).¹⁸

Discussion

We found that participating in either a medically tailored meals intervention or a nontailored food intervention for at least six months was associated with significant reductions in ED visits, compared with visits among similar participants who did not receive an intervention. These programs were also associated with lower use of other big-ticket health services. Sensitivity analyses conducted on the entire unmatched cohort were generally similar to the matched analyses,

and we did not see evidence that regression to the mean, use of other social supports, or a triggering event explained these findings.

The medically tailored meals and nontailored food interventions were used in different populations. In general, the former was used in a younger group with higher rates of disability, while the latter was used in an older group whose members were less likely to use English as their primary language. Information regarding these programs may be useful to policy makers and payers, particularly to the extent that the populations they are providing care for are similar to these groups. The medically tailored meals program may be particularly useful for those who are sicker and disabled. The nontailored food program may be a useful, and less expensive, way to improve health care use for people who are older but otherwise relatively healthy. For both programs, the overall reductions in ED visits are large enough to be clinically meaningful. Though we did not conduct a formal cost-effectiveness analysis in this study, the lower estimated spending suggests that these programs may offer savings to payers or at least be cost-neutral, while reducing unplanned and disruptive events such as ED visits and emergency transportation. Thus, the programs are likely to be beneficial for both patients and health care systems.

This study extends our knowledge of the effects of providing home-delivered meals on health outcomes. Few prior studies have examined the impact of medically tailored meals, but one study that did, in a different patient population, found cost savings associated with the meals.³⁰ With regard to the delivery of nontailored food, prior studies have suggested that home-delivered meals decrease nursing home admissions, but these studies were limited by their ecological design.^{31,32} Furthermore, a recent systematic review of home-delivered meal interventions found that most studies were cross-sectional and did not examine health care use or spending, as this study did.³³

This study suggests several directions for future work. Randomized evaluation for this type of intervention is feasible, and given our promising findings, that would be an important next step. It will also be important to confirm these findings in larger samples and in different settings, to evaluate the results' generalizability. Our results also support the overall approach of increasing the integration between health care and social services sectors.^{16,34,35}

This study had several key strengths. We used detailed assessment of pre-intervention data across several domains (clinical, claims-based, demographic, social, and geographical), which helped minimize confounding related to inter-

EXHIBIT 4

Estimated changes in utilization and spending based on matching using pre-intervention services (triggering events)

	Incidence rate ratio	(95% CI)
EMERGENCY DEPARTMENT VISITS		
Medically tailored meals program	0.36***	(0.24, 0.56)
Nontailored food program	0.54***	(0.42, 0.68)
INPATIENT ADMISSIONS		
Medically tailored meals program	0.41**	(0.19, 0.90)
Nontailored food program	0.81	(0.60, 1.10)
EMERGENCY TRANSPORTATION EVENTS		
Medically tailored meals program	0.18***	(0.08, 0.38)
Nontailored food program	0.52***	(0.39, 0.69)
Average monthly spending		
DIFFERENCE IN GROSS SPENDING		
Medically tailored meals program	-\$802***	(-1,363, -242)
Nontailored food program	-\$228**	(-380, -76)

SOURCE Authors' analysis of data from the Commonwealth Care Alliance. **NOTES** Spending is in inflation-adjusted 2016 dollars. Estimates from negative binomial models adjusted for the factors listed in the notes to exhibit 2. ***p* < 0.05 ****p* < 0.01

vention assignment. Our data were also longitudinal, and the participants were racially and ethnically diverse. Finally, the results were robust to several sensitivity analyses.

Conclusion

Home-delivered meals—in particular, those tailored to recipients' medical needs—show prom-

ise for helping curtail the use of selected costly health services in adults dually eligible for Medicare and Medicaid, a medically and socially complex population for whom effective interventions can be hard to come by. While further, preferably randomized, evaluations are needed, this study suggests that meal delivery may be an important way to improve the health of vulnerable patients. ■

An earlier version of this article was presented at the Root Cause Coalition National Summit, Louisville, Kentucky, October 9, 2017. This project was supported by a grant from the AARP Foundation. Seth Berkowitz's role in the research reported in this publication was also supported, in part, by the

National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health (Award No. K23DK109200). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Jean Terranova is an

employee of Community Servings. Caterina Hill, Todd Linsky, and Lori Tishler are employees of Commonwealth Care Alliance. Toyin Ajayi was an employee of Commonwealth Care Alliance during the study period.

NOTES

- 1 Adler NE, Newman K. Socioeconomic disparities in health: pathways and policies. *Health Aff (Millwood)*. 2002;21(2):60–76.
- 2 Berwick DM, Nolan TW, Whittington J. The Triple Aim: care, health, and cost. *Health Aff (Millwood)*. 2008;27(3):759–69.
- 3 Wilensky G. Addressing social issues affecting health to improve US health outcomes. *JAMA*. 2016;315(15):1552–3.
- 4 Coleman-Jensen A, Rabbitt MP, Gregory CA, Singh A. Household food security in the United States in 2015 [Internet]. Washington (DC): Department of Agriculture, Economic Research Service; 2016 Sep [cited 2018 Feb 7]. (Economic Research Report No. 215). Available from: <https://www.ers.usda.gov/webdocs/publications/79761/err-215.pdf>
- 5 Bhargava V, Lee JS. Food insecurity and health care utilization among older adults in the United States. *J Nutr Gerontol Geriatr*. 2016;35(3):177–92.
- 6 Heflin C, Hodges L, Mueser P. Supplemental Nutrition Assistance Program benefits and emergency room visits for hypoglycaemia. *Public Health Nutr*. 2017;20(7):1314–21.
- 7 Berkowitz SA, Meigs JB, DeWalt D, Seligman HK, Barnard LS, Bright OJ, et al. Material need insecurities, control of diabetes mellitus, and use of health care resources: results of the Measuring Economic Insecurity in Diabetes study. *JAMA Intern Med*. 2015;175(2):257–65.
- 8 Kushel MB, Gupta R, Gee L, Haas JS. Housing instability and food insecurity as barriers to health care among low-income Americans. *J Gen Intern Med*. 2006;21(1):71–7.
- 9 Berkowitz SA, Baggett TP, Wexler DJ, Huskey KW, Wee CC. Food insecurity and metabolic control among U.S. adults with diabetes. *Diabetes Care*. 2013;36(10):3093–9.
- 10 Berkowitz SA, Berkowitz TSZ, Meigs JB, Wexler DJ. Trends in food insecurity for adults with cardiometabolic disease in the United States: 2005–2012. *PLoS One*. 2017;12(6):e0179172.
- 11 Leung CW, Epel ES, Willett WC, Rimm EB, Laraia BA. Household food insecurity is positively associated with depression among low-income Supplemental Nutrition Assistance Program participants and income-eligible nonparticipants. *J Nutr*. 2015;145(3):622–7.
- 12 Gundersen C, Ziliak JP. Food insecurity and health outcomes. *Health Aff (Millwood)*. 2015;34(11):1830–9.
- 13 Seligman HK, Schillinger D. Hunger and socioeconomic disparities in chronic disease. *N Engl J Med*. 2010;363(1):6–9.
- 14 Berkowitz SA, Basu S, Meigs JB, Seligman HK. Food insecurity and health care expenditures in the United States, 2011–2013. *Health Serv Res*. 2017 Jun 13. [Epub ahead of print].
- 15 Bynum JPW, Austin A, Carmichael D, Meara E. High-cost dual eligibles' service use demonstrates the need for supportive and palliative models of care. *Health Aff (Millwood)*. 2017;36(7):1309–17.
- 16 Alley DE, Asomugha CN, Conway PH, Sanghavi DM. Accountable Health Communities—addressing social needs through Medicare and Medicaid. *N Engl J Med*. 2016;374(1):8–11.
- 17 Commonwealth Care Alliance. About our unique healthcare organization [Internet]. Boston (MA): CCA; c 2018 [cited 2018 Feb 7]. Available from: <http://www.commonwealthcarealliance.org/about>
- 18 To access the appendix, click on the Details tab of the article online.
- 19 CMS.gov. Risk adjustment [Internet]. Baltimore (MD): Centers for Medicare and Medicaid Services; [last modified 2017 Jan 3; cited 2018 Feb 7]. Available from: <https://www.cms.gov/Medicare/Health-Plans/MedicareAdvSpecRateStats/Risk-Adjustors.html>
- 20 Gagne JJ, Glynn RJ, Avorn J, Levin R, Schneeweiss S. A combined comorbidity score predicted mortality in elderly patients better than existing scores. *J Clin Epidemiol*. 2011;64(7):749–59.
- 21 Wong ES, Rosland AM, Fihn SD, Nelson KM. Patient-centered medical home implementation in the Veterans Health Administration and primary care use: differences by patient comorbidity burden. *J Gen Intern Med*. 2016;31(12):1467–74.
- 22 Census Bureau. American Community Survey: data tables and tools [Internet]. Washington (DC): Census Bureau; [cited 2018 Feb 7]. Available from: <https://www.census.gov/acs/www/data/data-tables-and-tools/>
- 23 Dartmouth Atlas of Health Care. Downloads [Internet]. Lebanon (NH): Trustees of Dartmouth College; c 2018 [cited 2018 Feb 7]. Available from: <http://www.dartmouthatlas.org/tools/downloads.aspx>
- 24 Stevens GA, King G, Shibuya K. Deaths from heart failure: using coarsened exact matching to correct cause-of-death statistics. *Popul Health Metr*. 2010;8:6.
- 25 Iacus SM, King G, Porro G. Multivariate matching methods that are monotonic imbalance bounding. *J Am Stat Assoc*. 2011;106(493):345–61.
- 26 Iacus SM, King G, Porro G. Causal inference without balance checking: coarsened exact matching. *Polit Anal*. 2012;20(1):1–24.
- 27 Iacus S, King G, Porro G. CEM: coarsened exact matching software [Internet]. Cambridge (MA): Har-

- vard Institute for Quantitative Social Science; [cited 2018 Feb 7]. Available from: <https://gking.harvard.edu/cem>
- 28** Barnett AG, van der Pols JC, Dobson AJ. Regression to the mean: what it is and how to deal with it. *Int J Epidemiol.* 2005;34(1):215–20.
- 29** Manning WG, Mullahy J. Estimating log models: to transform or not to transform? *J Health Econ.* 2001; 20(4):461–94.
- 30** Gurley J, Rand K, Daugherty S, Dinger C, Schmeling J, Laverty N. Examining health care costs among MANNA clients and a comparison group. *J Prim Care Community Health.* 2013;4(4):311–7.
- 31** Thomas KS, Mor V. Providing more home-delivered meals is one way to keep older adults with low care needs out of nursing homes. *Health Aff (Millwood).* 2013;32(10): 1796–802.
- 32** Thomas KS, Mor V. The relationship between Older Americans Act Title III state expenditures and prevalence of low-care nursing home residents. *Health Serv Res.* 2013;48(3): 1215–26.
- 33** Campbell AD, Godfryd A, Buys DR, Locher JL. Does participation in home-delivered meals programs improve outcomes for older adults? Results of a systematic review. *J Nutr Gerontol Geriatr.* 2015;34(2): 124–67.
- 34** Barnidge E, Stenmark S, Seligman H. Clinic-to-community models to address food insecurity. *JAMA Pediatr.* 2017;171(6):507–8.
- 35** Berkowitz SA, Hulberg AC, Standish S, Reznor G, Atlas SJ. Addressing unmet basic resource needs as part of chronic cardiometabolic disease management. *JAMA Intern Med.* 2017;177(2):244–52.